arXiv:1301.2873v1 [hep-ph] 14 Jan 2013

Net quark number probability distribution near the chiral crossover transition

Kenji Morita,[1, 2, *] Bengt Friman,[3] Krzysztof Redlich,[4, 2] and Vladimir Skokov[5]

[1] *Yukawa Institute for Theoretical Physics, Kyoto University, Kyoto 606-8502, Japan*
[2] *Extreme Matter Institute EMMI, GSI, Planckstr. 1, D-64291 Darmstadt, Germany*
[3] *GSI, Helmholzzentrum für Schwerionenforschung, Planckstr. 1, D-64291 Darmstadt, Germany*
[4] *Institute of Theoretical Physics, University of Wroclaw, PL-50204 Wrocław, Poland*
[5] *Physics Department, Brookhaven National Laboratory, Upton, NY 11973, USA*

(Dated: January 15, 2013)

We investigate properties of the probability distribution of the net quark number near the chiral transition in the quark-meson model. The calculations are performed within the functional renormalization group method and under mean-field approximation. We show, that there is a substantial influence of the remnant of the chiral phase transition on the properties of the probability distribution. In particular, at physical pion mass, the presence of mesonic fluctuations modifies the distribution from the mean-field one, as well as, from the Skellam distribution which is set as the reference for a non critical behavior. An intrinsic feature of the net quark number probability distribution is, that in the vicinity of the chiral crossover transition it is narrower than the corresponding Skellam function. We also discuss scaling properties of the probability distribution and the resulting cumulants.

I. INTRODUCTION

The structure of the QCD phase diagram is one of the fundamental problems addressed in both, theoretical and experimental studies [1, 2]. At finite chemical potential, there have been predictions for the existence of the QCD critical point (CP), both within effective chiral models [3] and some lattice QCD calculations [4]. Despite extensive studies, however, the exact location of the CP and even its existence have not been established yet.

So far, the lattice QCD simulation at physical quark masses has shown, that at vanishing baryon density the transition from hadrons to a quark matter is crossover, in which, no associated singularities exist in thermodynamic observables [5]. Therefore, possible experimental signatures of the phase transition could be washed out. However, the chiral crossover transition in the QCD medium was conjectured [6] to appear as remnants of the second order phase transition belonging to the $O(4)$ universality class. Indeed, recent lattice QCD calculations have confirmed the $O(4)$ scaling behavior of relevant physical observables at the physical quark masses [7, 8]. This result has opened new opportunities to verify the QCD phase boundary experimentally by measuring fluctuations of conserved charges [9–13].

The fluctuations of conserved charges, especially of the net baryon number and the electric charge, were proposed to search for the conjectured CP [9, 14–18]. However, in view of remnants of the $O(4)$ criticality and by virtue of the proximity of the chiral crossover to the freeze-out line in heavy ion collisions, fluctuations of the conserved charges were also proposed as diagnostic tools of the QCD phase boundary outside the CP [9, 10, 16, 19–21]. It has been shown, that at chiral crossover transition, there should appear a non-trivial structure of the higher order cumulants of the net baryon number and electric charge due to $O(4)$ criticality [20, 21]. The theoretically expected modifications of different cumulants due to the chiral dynamics can be partly observed in recent data of STAR Collaboration [22, 23]. The cumulants have also been studied in the first principle lattice QCD approach [24–26], as well as, in effective chiral models [19–21, 27–33].

Fluctuations of conserved charges can be linked to the corresponding probability distribution. Thus, any critical behavior in cumulants of conserved charges should be also reflected in their probability distributions. Recently, we have examined the effects of the phase transition on the net baryon number probability distribution based on the Laudau theory of the phase transition [34]. We have argued that there is a substantial influence of criticality on the probability distribution caused by the existence of the singularities of the thermodynamic potential in the complex chemical potential plane. In the present work, we extend our previous studies to more realistic model. We consider the two-flavor quark-meson model which is a low energy effective approach to the chiral properties of QCD. We include consistently the critical fluctuations, which were introduced *ad hoc* in Ref. [34], by means of the functional renormalization group (FRG) method [35–37]. Although we neglect coupling to the Polyakov loop, which is important to implement the statistical confinement, the $O(4)$ scaling behavior and thermodynamics near the critical point is well described by this model within the FRG method [38, 39]. Clearly, some properties related with confinement in QCD, like e.g. relevant behavior of kurtosis near the chiral transition [16, 31] cannot be addressed in this approach. Therefore, we restrict our studies entirely to effects of the chiral transition on the net quark number probability distribution.

This paper is organized as follows: in the next section, we introduce the quark-meson model and its thermody-

---

[*] Electronic address: kmorita@yukawa.kyoto-u.ac.jp

namic properties. In Sec. III, we present results on the probability distribution and cumulants of the net quark number. Section IV is devoted to the concluding remarks.

II. THE THERMODYNAMIC POTENTIAL IN THE QUARK-MESON MODEL

We employ the quark-meson model to explore the influence of the chiral phase transition on the probability distribution of the net quark-number.

The quark-meson model is an effective realization of the low energy QCD in which the meson field $\phi = (\sigma, \vec{\pi})$, in the $O(4)$ representation, is coupled to quark fields through the covariant derivative. The Lagrangian density is then given by

$$\mathcal{L} = \bar{q}[i\gamma_\mu \partial^\mu - g(\sigma + i\gamma_5 \vec{\tau}\cdot\vec{\pi})]q + \frac{1}{2}(\partial_\mu \sigma)^2 + \frac{1}{2}(\partial_\mu \vec{\pi})^2 - U(\sigma, \vec{\pi}), \tag{1}$$

where $U(\sigma)$ denotes the mesonic potential,

$$U(\sigma, \vec{\pi}) = \frac{1}{2}m^2\phi^2 + \frac{\lambda}{4}\phi^4 - h\sigma. \tag{2}$$

For $m^2 < 0$ and $h = 0$, the $O(4)$ symmetry of the potential is spontaneously broken to $O(3)$, resulting in a non-vanishing value of the vacuum scalar condensate $\langle\sigma\rangle$. The last term, $h = f_\pi m_\pi^2$, breaks explicitly the chiral symmetry and yields the nonzero pion mass.

To describe thermodynamics of the quark-meson model we calculate thermodynamic potential within the FRG approach as introduced in Ref. [38]. Following [35], we consider a scale dependent effective action in the local potential approximation. We neglect the wave function renormalization and the flow of the Yukawa coupling. Then, using the so-called optimized cutoff functions, the evolution equation for the scale dependent thermodynamic potential density with the reduced field varible $\rho = (\sigma^2 + \vec{\pi}^2)/2$ is found as [38],

$$\partial_k \Omega_k(\rho) = \frac{k^4}{12\pi^2}\left[\frac{3}{E_\pi}\left\{1 + 2n_B(E_\pi)\right\} + \frac{1}{E_\sigma}\left\{1 + 2n_B(E_\sigma)\right\} - \frac{2\nu_q}{E_q}\left\{1 - n_F(E_q^+) - n_F(E_q^-)\right\}\right], \tag{3}$$

where n_B and n_F are the Bose and the Fermi distribution functions, respectively, $\nu_q = 2N_c N_f = 12$ stands for the quark degeneracy. The single particle energies of pion, sigma meson and quark/antiquark read,

$$E_\pi = \sqrt{k^2 + \bar{\Omega}_k'}, \quad E_\sigma = \sqrt{k^2 + \bar{\Omega}_k' + 2\rho\bar{\Omega}_k''}, \quad E_q^\pm = \sqrt{k^2 + 2g^2\rho} \pm \mu, \tag{4}$$

where $\bar{\Omega}_k'$ and $\bar{\Omega}_k''$ denote the first and the second derivatives of $\bar{\Omega}_k = \Omega_k + h\sqrt{2\rho_k}$, with respect to ρ. The flow equation (3) is solved within the Taylor expansion method. Expanding the potential up to the third order in ρ around the potential minimum ρ_k,

$$\Omega_k(\rho) = \sum_{n=0}^{3} \frac{a_n(k)}{n!}(\rho - \rho_k)^n, \tag{5}$$

and using Eq. (3), one gets the flow equations for $a_n(k)$ and ρ_k, which can be solved numerically starting at a cutoff scale $\Lambda = 1.0$ GeV [38]. We eliminate a_1 by the scale independent relation $h = a_1(k)\sqrt{2\rho_k}$. Thus, there are four initial parameters in the flow equations to be fixed at the cutoff scale $k = \Lambda$. The $a_0(\Lambda)$ gives is a constant shift of Ω. The finite momentum cutoff implies a non physical high temperature behavior of thermodynamic quantities due to the lack of high momentum contribution beyond the cutoff scale [31, 40]. Thus, following the prescription introduced in Ref. [40], we include the high momentum contribution by considering the flow equation for the non-interacting massless quarks and gluons,

$$\partial_k \Omega_k^\Lambda(T,\mu) = \frac{k^3}{12\pi^2}\{2(N_c^2 - 1)[1 + 2n_B(k)] - \nu_q[1 - n_F(k^+) - n_F(k^-)]\}. \tag{6}$$

Integrating the flow equation (6) from $k = \infty$ to $k = \Lambda$, we obtain $\Omega^\Lambda(T,\mu)$ which is used for $a_0(\Lambda)$. If one puts $a_3(\Lambda) = 0$ then the remaining two parameters, the running minimum $\rho_{k=\Lambda}$ and $a_2(\Lambda)$ are fixed such that to reproduce the pion $m_\pi = 135$ MeV and the sigma $m_\sigma = 640$ MeV masses. The strength of the Yukawa coupling, $g = 3.2$, is fixed by the constituent quark mass, $M_q(T = \mu = 0) = g\sigma_{k=0}(T = \mu = 0) = 300$ MeV with $\sigma_{k=0}(T = \mu = 0) = f_\pi = 93$ MeV. The full thermodynamic potential density of the quark-meson model $\Omega(T,\mu)$, which includes thermal and quantum mesonic fluctuations, is obtained from $\Omega(T,\mu) = \lim_{k\to 0}\Omega_k$, where $\Omega_{k=0}$ is the solution of the flow equation (3).

If one ignores the flow of the mesonic part in the evolution equation (3), the resulting effective potential corresponds to that obtained in the mean-field approximation with a finite cutoff. The fermionic vacuum fluctuation term, included in the mean-field (MF) potential, is necessary to reproduce the correct order of the phase transition at vanishing μ. This vacuum term has to be renormalized to remove the artifact of the cutoff dependence [19]. Then, in the mean-field approximation, the thermodynamic potential is obtained as

$$\Omega_{\rm MF}(\langle\sigma\rangle;T,\mu) = U(\langle\sigma\rangle,\vec{\pi}=0) - \frac{\nu_q}{16\pi^2}M_q^4\ln\left(\frac{M_q}{M}\right)$$
$$-\nu_q T\int\frac{d^3p}{(2\pi)^3}\left[\ln(1+e^{-(E_q-\mu)/T}) + \ln(1+e^{-(E_q+\mu)/T})\right], \tag{7}$$

where $M_q = g\langle\sigma\rangle$ and the expectation value $\langle\sigma\rangle$ is determined as the solution of the gap equation $\partial\Omega_{\rm MF}/\partial\langle\sigma\rangle = 0$.

III. PROBABILITY DISTRIBUTION OF THE NET QUARK NUMBER

The thermodynamical potentials for the quark-meson model, derived in the previous section, can be used to calculate the influence of the chiral phase transition, or its remnant, on different observables. In the following we concentrate on properties of the probability distribution $P(N)$ and the resulting cumulants $c_n(T,\mu)$ of the net quark number N.

A. General properties of $P(N)$

We consider a thermodynamic system described by the grand canonical ensemble at temperature T in a subvolume V. We introduce the chemical potential μ which is connected to the average net charge conservation. For the net quark number $N = N_q - N_{\bar{q}}$, the probability distribution $P(N)$ to find the net charge N in the volume V is given by

$$P(N;T,\mu,V) = \frac{Z(T,V,N)}{\mathcal{Z}(T,\mu,V)}e^{\mu N/T}, \tag{8}$$

where $Z(T,V,N)$ is the canonical and $\mathcal{Z}(T,\mu,V)$ the grand-canonical partition function. Owing to the condition, $\sum_{N=-\infty}^{\infty}P(N) = 1$, the grand canonical partition function in Eq. (8) plays a role of the normalization. Thus, all essential information on $P(N)$ properties is contained in the canonical partition function. The $Z(T,V,N)$ is connected with the grand canonical partition function through the fugacity expansion,

$$\mathcal{Z}(T,V,\lambda) = \sum_N \lambda^N Z(T,V,N), \tag{9}$$

where $\lambda = e^{\beta\mu}$. Consequently, the canonical partition function is obtained as the coefficient of this Laurent series,

$$Z(T,V,N) = \frac{1}{2\pi i}\oint_C d\lambda\frac{\mathcal{Z}(T,V,\lambda)}{\lambda^{N+1}}. \tag{10}$$

Thus, to find the canonical partition function one needs to know the analytic structure of $\mathcal{Z}(T,V,\lambda)$ in the complex λ plane to choose an appropriate integration contour.[1] In the chiral effective models, the general structure of the singularities associated with the chiral phase transition has been discussed in Refs. [41, 42]. In the broken phase, $T < T_c(\mu)$, there is no singularity on the unit circle $\lambda = e^{i\theta}$ with $0 \le \theta < 2\pi$. Consequently, the canonical partition function can be represented as [43, 44],

$$Z(T,V,N) = \frac{1}{2\pi}\int_0^{2\pi}d\theta e^{-i\theta N}\mathcal{Z}(T,V,\theta), \tag{11}$$

where $\theta = \mu_I/T$ is the imaginary chemical potential.

The above equation links the grand canonical partition function at imaginary chemical potential to thermodynamics at a finite and fixed charge. The QCD partition function at imaginary chemical potential exhibits the Roberge-Weiss periodicity [45], $\mathcal{Z}(T,V,\theta+2\pi/3) = \mathcal{Z}(T,V,\theta)$, which implies the quantization of the net quark number by a multiple of three [46–48]. Owing to the lack of confinement, the quark-meson model has the periodicity 2π in θ, thus here N is an integer.

In the present work, following Ref. [34], we will use Eq. (11) to calculate the canonical partition function and the corresponding probability distribution. The thermodynamic potential density $\Omega = -(T/V)\log\mathcal{Z}$ will be calculated in the quark-meson model within the FRG approach and under the mean-field approximation. We also assume that the thermodynamic potential density is well approximated by its thermodynamic limit. Such an approximation, also employed in Ref. [34], is justified for $|N| \ll N^*$, where N^* is a characteristic of the partition function of a finite system in the Yang-Lee theory of phase transitions [49].

Because of an oscillating structure of the integrand, the numerical integration of Eq. (11) has a limited applicability at large $|N|$. We found, however, that the numerical integration works up to the value of $|N|$ at which $P(N;\mu=0) \sim 10^{-12}$, independent of the volume, temperature and other parameters. Such numerical accuracy is sufficient to study the influence of the chiral criticality on the properties of the net quark number probability distribution.

B. The net quark number distribution near the chiral phase transition

In the chiral limit and at vanishing and moderate values of μ, the quark-meson model exhibits a second order phase transition which belongs to the $O(4)$ universality class. For a physical pion mass, the chiral symmetry is explicitly broken. Nevertheless, there is a remnant of the $O(4)$ criticality in different observables [9]. In particular,

---

[1] In a finite system, the grand partition function $\mathcal{Z}$ has Yang-Lee zeroes at complex λ, which turn into cuts in the thermodynamic limit.



FIG. 1: Probability distributions of the net quark number just below the pseudocritical temperature T_{pc} in the FRG approach (left) and in the mean-field (MF) approximation (right), compared with the corresponding Skellam distributions. The dots are the model results for different volumes while the lines stand for the Skellam distribution.



FIG. 2: Probability distribution for different temperatures in the FRG approach (left) and in the mean-field (MF) approximation (right). The dots are the model results while the lines stand for the corresponding Skellam distribution.

such a remnant should be also present in the probability distribution of the net quark number distribution which is governed by the properties of the thermodynamic potential. Since the $P(N)$ is calculated at finite volume, to see the influence of the long range critical correlations, the probability distribution should be calculated for a sufficiently large V and N and also near the pseudocritical temperature T_{pc}.

To disentangle the influence of the chiral criticality on $P(N)$, one needs to establish the reference. At low temperatures, $T \ll T_{pc}$, the thermodynamic potential is well approximated as a quasi-ideal quark gas with dynamically generated mass which is $T-$dependent. Consequently, at any fixed T and V, the natural reference for $P(N)$ is the Skellam distribution which reflects the probability distribution calculated from the non-singular part, an ideal gas approximation, of the thermodynamic potential [10]. The Skellam distribution, in this case [10], is quantified by the mean number of quarks $b = \langle N_q \rangle$ and antiquarks $\bar{b} = \langle N_{\bar{q}} \rangle$ as,

$$P(N) = \left(\frac{b}{\bar{b}}\right)^{N/2} I_N(2\sqrt{b\bar{b}})e^{-(b+\bar{b})}, \qquad (12)$$

where $I_N(x)$ is the modified Bessel function.

In the quark-meson model, b and $\bar{b}$ are calculated for an ideal gas of constituent quarks with dynamically generated mass

$$b = \frac{\nu_q V}{2\pi^2}\int_0^\infty dk k^2 n_F(E_k; T, \mu), \qquad (13)$$

where n_F is the Fermi distribution function and $E_k = \sqrt{k^2 + M_q^2}$ is the quasi-quarks energy. In the MF approximation $M_q = g\langle\sigma\rangle$, while in the FRG approach we apply the scale dependent quark mass $M_{q,k} = g\sigma_k$ which for $k \leq \Lambda$ is obtained from the flow equation (3). For $\Lambda < k < \infty$ the quarks are assumed to be massless and non-interacting.

Figure 1 shows the probability distributions of the net quark number obtained in the quark-meson model, both



FIG. 3: Probability distributions in FRG (red, circles), Skellam (blue, dashed), and free fermi gas (green, solid).

in the FRG approach (left) and under the MF approximation (right). The calculations were done at vanishing chemical potential in the vicinity of the pseudocritical temperature T_{pc}. The values of T_{pc} were determined by the peak position of the chiral susceptibility which in the FRG and MF cases correspond to 214 MeV and 190 MeV, respectively. $P(N)$ is seen in Fig. 1 to be strongly changing with the volume of the system. With increasing V the distribution is broader as expected, since the variance of the distribution is linearly dependent on the volume. There is also a strong change of the distribution when increasing distance from the pseudocritical point. As seen in Fig. 2, the distribution is narrower for smaller values of T/T_{pc} at a fixed volume. This is directly connected to a decreasing mean number of quarks in the system.



FIG. 4: Cumulants c_n $(n = 2, 4, 6)$ of the net charge with the mean $b = \bar{b} = 6.0$ calculated for the Skellam distribution from Eqs. (16)-(19) for different values of $N_{\rm max}$.

In comparison, both the MF and the FRG distributions are narrower than the Skellam counterparts. The deviation becomes larger when the volume increases and the mesonic fluctuations are included, as seen in Fig. 1. The effect of the mesonic fluctuations is more clearly visible when temperature is varied. In Fig. 2, we plot $P(N)$ for three different temperatures below T_{pc}. Except for the highest temperature, the $P(N)$ in the MF calculations coincides with the Skellam distribution, while it deviates in the FRG case. As temperature approaches to T_{pc}, the difference between the Skellam and the model results becomes larger, indicating an increasing influence of remnants of the $O(4)$ criticality due to the chiral dynamics.

While the shrinking of the $P(N)$ in the FRG calculations from the Skellam distribution could be expected from the previous studies of the Laudau theory in Ref. [34], one may ask why such behavior is also seen in the present MF approach. As discussed in [34], the shrinking of $P(N)$ originates from the branch point in a complex chemical potential plane and its associated cut, near which, the two saddle points are located. Although the σ^4 model used in Ref. [34] does not have a cut in thermodynamic potential but in the order parameter, in general, however, such a cut exists also in the thermodynamic potential under the MF approximation of the quark-meson model, leading to the observed shrinking of $P(N)$ below the Skellam distribution.

The Skellam distribution (12), corresponds to a non-interacting gas of classical particles. In general, effects of quantum statistics should be taken into account when the mass of quark is small relative to the temperature. In Eq. (13), we use the Fermi distribution function, thus the effect of quantum statistics is partly taken into account through the number of quarks and antiquarks. Nevertheless, the quantum statistics can also change the functional form of the probability distribution, as shown for the pionic electric charge fluctuations in Ref. [11]. In the quark-meson model within the FRG calculation one gets, $M_q \simeq 126$ MeV at $T/T_{pc} = 0.98$, the mass, which is too small to verify the Boltzmann approximation near T_{pc}. Therefore, to disentangle the criticality and the quantum statistics effect, we also calculate the probability distribution $P(N)$ for the free quark and anti-quark gas through Eq. (11) from the thermodynamic potential density

$$\Omega_{\rm fermi}(T,\theta) = -\frac{\nu_q T}{2\pi^2}\int_0^\infty dk k^2[\ln(1+e^{-E_k/T+i\theta}) + \ln(1+e^{-E_k/T-i\theta})] \qquad (14)$$

with the same E_k as used in Eq. (13). Note, that by taking the Boltzmann approximation in Eq. (14) one recovers the Skellam distribution (12) in which the baryon number is given by the Boltzmann distribution.

Figure 3 shows, that the probability distributions $P(N)$ for the free Fermi gas exhibits narrower width than the corresponding Skellam distribution, reflecting the influence of the quantum statistics. However, the $P(N)$ obtained in the quark-meson model within the FRG approach is still narrower than for the corresponding Fermi gas, indicating the effect of the $O(4)$ criticality.



FIG. 5: Cumulants c_n $(n = 2, 4, 6)$ calculated in the quark-meson model within the FRG approach from Eq. (15) (full line) and from Eqs. (16)-(19) (points) for different values of $N_{\max}$. Results of the corresponding Skellam distribution with sufficiently large $N_{\max}$ are also indicated by open circles and dashed lines.



FIG. 6: Cumulants c_n $(n = 2, 4, 6)$ calculated in the quark-meson model within the FRG approach from Eq. (15) (full line) and from Eqs. (16)-(19) (points) for different values of $N_{\max}$.

C. Cumulants of the net quark number

Cumulants, $c_n(T, \mu)$ of the net-quark number, are known to be excellent probe of the chiral phase transition and its remnant [9, 10, 16, 19–21]. In the statistical physics, c_n are quantified by the corresponding susceptibilities,

$$c_n(T, \mu) \equiv \frac{\partial^n [p(T, \mu)/T^4]}{\partial(\mu/T)^n}. \tag{15}$$

Thus, once the thermodynamic pressure $p(T, \mu, V) = (T/V) \ln \mathcal{Z}$ is calculated in the grand-canonical ensemble, the cumulants $c_n(T, \mu)$ can be obtained by taking derivatives of the thermodynamic pressure with respect to the chemical potential. Clearly, since the higher order cumulants are increasingly sensitive to the chiral phase transition, their calculations involve special numerical efforts [9, 24, 50]. However, once the probability distribution of the net charge multiplicity, Eq. (8), is known, the cumulants can be calculated directly from $P(N)$ in terms of powers of the central moments $\langle(\delta N)^k\rangle = \langle(N - \langle N\rangle)^k\rangle$. Indeed, at $\mu = 0$, the first three non-vanishing cumulants c_n $(n = 2, 4, 6)$ are expressed by the central moments as

$$c_2 = \frac{\langle(\delta N)^2\rangle}{VT^3}, \tag{16}$$

$$c_4 = \frac{\langle(\delta N)^4\rangle - 3\langle(\delta N)^2\rangle}{VT^3}, \tag{17}$$

$$\begin{aligned} c_6 = {} & \left[\langle(\delta N)^6\rangle - 15\langle(\delta N)^4\rangle\langle(\delta N)^2\rangle \right. \\ & \left. -10\langle(\delta N)^3\rangle + 30\langle(\delta N)^2\rangle\right]/(VT^3); \end{aligned} \tag{18}$$

where $\langle N^k\rangle$ are directly obtained from the probability distribution

$$\langle N^k\rangle = \sum_{N=-N_{\max}}^{N_{\max}} N^k P(N), \tag{19}$$

where in general, $N_{\max} = \infty$. In practice, however, the $N_{\max}$ is always finite. Consequently, the cumulants calculated from Eqs. (16)-(19) are only approximations of exact results obtained by taking derivatives of the thermodynamic pressure as in Eq. (15). To illustrate this property we calculate different cumulants of Skellem distribution for different $N_{\max}$ and compare those results with exact calculations. Figure 4 shows the ratio of different moments as a function of $N_{\max}$ for a fixed mean number of quarks. It is clear from this figure, that to reproduce the exact results for different cumulants, one needs to calculated $P(N)$ for sufficiently large $N_{\max}$, which is increasing with the order of the cumulant. Clearly, the value of $N_{\max}$ which is required to get the best description of different cumulants also varies with the mean charge number. In the actual model calculations, the mean quark number depends, in general, on the volume, the temperature and the chemical potential.

Figure 5 shows the temperature dependence of different cumulants calculated in the quark-meson model within the FRG approach from Eq. (15) and their approximate values obtained from Eqs. (16)-(19) for different $N_{\max}$. The cumulants obtained from the corresponding Skellam distribution are also shown in this figure for comparison.

The convergence properties of different cumulants to their exact values with $N_{\max}$, in the quark-meson model, are similar to that already seen in the Skellam distribution. The $N_{\max}$ needed to get a good approximation of the exact results are increasing with the order of cumulants, since they are more sensitive to the tail of the distribution. For the parameters used in Fig. 5, all cumulants, up to the sixth order, are well reconstructed from the probability distribution by choosing $N_{\max} \sim 90$. This also confirms consistency of the calculations of the canonical partition function.

At $\mu = 0$, the second and the fourth order cumulants are not influenced by the critical chiral dynamics, as they are finite even in the chiral limit. Their properties are entirely determined by the regular part of the partition function. The temperature dependence of c_2 and c_4, seen in Fig. 5, is due to the temperature dependence of the dynamical quark mass.

Contrary to c_2 and c_4, the temperature dependence of c_6 and its negative values near T_{pc}, seen in Fig. 5, has a universal structure. The characteristic shape of c_6 is typical for the $O(4)$ universality class, originated from the universal scaling function [9]. The c_6 obtained from the noncritical Skellam distribution, exhibits a very different behavior.

It is interesting to note, that already at moderate values of $N_{\max}$, the $O(4)$ shape structure of c_6 can be reproduced from the probability distribution. This is an interesting result for heavy ion experiments where one expects to observe the chiral criticality by reconstructing the higher order moments from a measured probability distribution of the net charge [22].

Deviations of c_n^A, obtained from the probability distribution (16)-(19), from their exact values c_n^E, obtained from Eq. (5), also depend on the volume of the system. However, there is an interesting approximate scaling relation (APS) which describes the relative deviation of c_n^A from c_n^E for different volumes. Figure 6 shows the relative difference, $R = (c_n^E - c_n^A)/c_n^E$ for $n = 2, 4, 6$, as a function of $N_{\max}$ normalized by $\sqrt{V}$. There is a clear APS of c_2 and those of higher order cumulants for sufficiently large $N_{\max}/\sqrt{V}$. Since for fixed T the mean number of quarks N_q scales linearly with the volume, one can replace $\sqrt{V}$ by $\sqrt{N_q}$ to preserve the observed approximate scaling properties.



FIG. 7: Rescaling of the net quark number probability distribution at fixed T with volume in the quark-meson model within the FRG approach and for the corresponding Skellam distribution.

The moments of the net quark number are linked to the corresponding probability distribution through Eq. (19). Thus, the APS of cumulants, seen in Fig. 6, should be also reflected in the properties of the net quark number probability distribution. Figure 7 shows, that for fixed T the rescaled probability distribution $\sqrt{V}P(N)$ indeed scales approximately with $N/\sqrt{V}$. The APS is valid for $P(N)$ obtained in the quark-meson model within the FRG approach, as well as, for the corresponding Skellam distribution. Therefore, the strong variation of $P(N)$ with V, seen in Fig. 1, can be essentially reduced if plotted as functions of the rescaled variable, $N/\sqrt{V}$. The results in Fig. 7 also confirm, that shrinking of the probability distribution of conserved charge, relative to the Skellam distribution, due to the $O(4)$ chiral dynamics is almost independent of the system volume.

Figure 8 shows probability distributions rescaled with the standard deviation of the distribution, $\sigma = \sqrt{\langle(\delta N)^2\rangle} = \sqrt{c_2 V T^3}$. Probability distributions obtained from the model calculation and the corresponding Skellam functions are hardly distinguishable on the logarithmic plot. They are also very different from the Gaussian distribution. The above results imply, that the global shape of the probability distribution with $O(4)$ criticality is not significantly different from the Skellam distribution, in spite of a very different behavior of the higher order cumulants. However, when approaching to the critical point, there is a clear deviation from the Skellam distribution even after rescaling.

Figure 9 demonstrates the violation of the "scaling behavior" close to the second order phase transition, in which the pion mass was reduced according to $h = 10^{-8} f_\pi m_\pi^2$. There is clear anomalous behavior in the tail of the distribution, which can be interpreted as the realization of the oscillation due to the saddle points dis-



FIG. 8: Rescaling of the probability distribution with the standard deviation $\sigma = \sqrt{\langle(\delta N)^2\rangle}$ for different volumes, for the FRG distribution in the quark-meson model and for the corresponding Skellam distribution. The standard normal distribution $(2\pi)^{-1/2}\exp(-x^2/2)$ is also plotted by the dashed line.



FIG. 9: Probability distribution of the quark-meson model in the chiral limit rescaled with the standard deviation for different volumes.

cussed in Ref. [34]. The deviation becomes more significant at smaller volumes, as expected.

IV. CONCLUDING REMARKS

We have studied the properties of the probability distribution $P(N)$ of the net quark number in the presence of the critical chiral dynamics governed by the $O(4)$ universality class in terms of the quark-meson model. Our calculations of $P(N)$ were done within the Functional Renormalization Group (FRG) approach which preserves the $O(4)$ scaling of relevant physical observables.

The main objectives of this paper was to verify the influence of the remnant of the chiral phase transition, expected at the physical pion mass, on the net quark probability distribution. We have shown that $P(N)$ is a sensitive observable to criticality by comparing the FRG results to the mean-field (MF) calculations. The thermal and quantum mesonic fluctuations, which are present in the FRG approach, imply modifications to the distribution obtained under the MF dynamics.

We have shown, that near the chiral crossover transition, the probability distribution of the net quark number is narrower than the Skellam distribution, which was used as the reference for the non-critical distribution at the corresponding temperature. Such narrowing of the probability distribution is dominated by the $O(4)$ criticality. In the quark meson model, however, it can also appear due to the quantum statistic effect as a consequence of a decreasing quark mass near the chiral transition. In the QCD medium, the masses of baryons are not directly effected by the critical chiral dynamics. Therefore, the narrowing of the net baryon number or the net proton number probability distribution measured in heavy ion collisions can be directly linked to the $O(4)$ criticality.

We have found an approximate scaling properties of the probability distribution and the net charge cumulants for different variables and thermal parameters. We have shown, that there is approximate scaling of $P(N)$ with the volume of the system, implying that the observed shrinking of $P(N)$ near the chiral transition is very weekly volume dependent. We expect, that our results are of importance in heavy ion collisions, where by measuring the net baryon number probability distribution and related moments, one expects to probe the QCD phase boundary experimentally. The phenomenological implications of our results will be presented elsewhere [51].

Acknowledgments

K.M. was supported by Yukawa International Program for Quark-Hadron Sciences at Kyoto University and by the Grant-in-Aid for Scientific Research from JSPS No.24540271. K.R. acknowledges partial support of the Polish Ministry of National Education (MEN). The research of V.S. is supported under Contract No. DE-AC02-98CH10886 with the U. S. Department of Energy. We acknowledge stimulating discussions with P. Braun-Munzinger, F. Karsch and Nu Xu.

---

[1] B. Friman, C. Höhne, J. Knoll, S. Leupold, J. Randrup, R. Rapp, and P. Senger, Lect. Note. Phys. **814**, 1 (2011).
[2] K. Fukushima and T. Hatsuda, Rep. Prog. Phys. **74**, 014001 (2011).
[3] M. Asakawa and K. Yazaki, Nucl. Phys. **A504**, 668 (1989).
[4] Z. Fodor and S. D. Katz, JHEP **0203**, 014 (2002); C. Schmidt, C. R. Allton, S. Ejiri, S. J. Hands, O. Kaczmarek, F. Karsch and E. Laermann, Nucl. Phys. Proc. Suppl. **119**, 517 (2003); R. V. Gavai, Nucl. Phys. A **862**, 104 (2011).
[5] Y. Aoki, G. Endrödi, Z. Fodor, S. D. Katz, and K. K. Szabó, Nature **443**, 675 (2006).
[6] R. D. Pisarski and F. Wilczek, Phys. Rev. D **29**, 338 (1984).
[7] S. Ejiri, F. Karsch, E. Laermann, C. Miao, S. Mukherjee, P. Petreczky, C. Schmidt, W. Soeldner, and W. Unger, Phys. Rev. D **80**, 094505 (2009).
[8] A. Bazavov, T. Bhattacharya, M. Cheng, C. DeTar, H. T. Ding, S. Gottlieb, R. Gupta and P. Hegde *et al.*, Phys. Rev. D **85**, 054503 (2012) [arXiv:1111.1710 [hep-lat]].
[9] F. Karsch and K. Redlich, Phys. Lett. B **695**, 136 (2011).
[10] P. Braun-Munzinger, B. Friman, F. Karsch, K. Redlich, and V. Skokov, Phys. Rev. C **84**, 064911 (2011).
[11] P. Braun-Munzinger, B. Friman, F. Karsch, K. Redlich, and V. Skokov, Nucl. Phys. **A880**, 48 (2012).
[12] O. Kaczmarek, F. Karsch, E. Laermann, C. Miao, S. Mukherjee, P. Petreczky, C. Schmidt, W. Soeldner, and W. Unger, Phys. Rev. D **83**, 014504 (2011).
[13] A. Bazavov *et al.* [HotQCD Collaboration], Phys. Rev. D **86**, 034509 (2012) [arXiv:1203.0784 [hep-lat]].
[14] Y. Hatta and M. A. Stephanov, Phys. Rev. Lett. **91**, 102003 (2003).
[15] M. Stephanov, K. Rajagopal, and E. Shuryak, Phys. Rev. Lett. **81**, 4816 (1998).
[16] S. Ejiri, F. Karsch, and K. Redlich, Phys. Lett. B **633**, 275 (2006).
[17] M. A. Stephanov, Phys. Rev. Lett. **102**, 032301 (2009).
[18] M. A. Stephanov, Phys. Rev. Lett. **107**, 052301 (2011).
[19] V. Skokov, B. Friman, E. Nakano, K. Redlich, and B.-J. Schaefer, Phys. Rev. D **82**, 034029 (2010).
[20] B. Friman, F. Karsch, K. Redlich, and V. Skokov, Eur. Phys. J. C **71**, 1694 (2011).
[21] V. Skokov, B. Friman, and K. Redlich, Phys. Rev. C **83**, 054904 (2011).
[22] M. M. Aggarwal et al. (STAR Collaboration), Phys. Rev. Lett. **105**, 022302 (2010).
[23] X. Luo (STAR Collaboration), talk given at Quark Matter 2012 International Conference, August 13-18, 2012, Washington D.C.
[24] C. R. Allton, M. Döring, S. Ejiri, S. J. Hands, O. Kaczmarek, F. Karsch, E. Laermann, and K. Redlich, Phys. Rev. D **71**, 054508 (2005).
[25] A. Bazavov et al. (HotQCD Collaboration), Phys. Rev. D **86**, 034509 (2012).
[26] S. Borsanyi, Z. Fodor, S. D. Katz, S. Krieg, C. Ratti, and K. Szabo (Wuppertal-Budapest Collaboration), JHEP **1201**, 138 (2012).
[27] K. Fukushima, Phys. Lett. B **591**, 277 (2004).
[28] C. Sasaki, B. Friman, and K. Redlich, Phys. Rev. D **75**, 054026 (2007).
[29] C. Sasaki, B. Friman, and K. Redlich, Phys. Rev. D **75**, 074013 (2007).
[30] B. Stokic, B. Friman, and K. Redlich, Phys. Lett. B **673**, 192 (2009).
[31] V. Skokov, B. Stokic, B. Friman, and K. Redlich, Phys. Rev. C **82**, 015206 (2010).
[32] M. Asakawa, S. Ejiri, and M. Kitazawa, Phys. Rev. Lett. **103**, 262301 (2009).
[33] T. K. Herbst, J. M. Pawlowski, and B. J. Schaefer, Phys. Lett. B **696**, 58 (2011).
[34] K. Morita, V. Skokov, B. Friman, and K. Redlich, arXiv:1211.4703 [hep-ph].
[35] C. Wetterich, Phys. Lett. B **301**, 90 (1993).
[36] J. Berges, N. Tetradis, and C. Wetterich, Phys. Rept **363**, 223 (2002).
[37] B. Delamotte, cond-mat/0702365.
[38] B. Stokić, B. Friman, and K. Redlich, Eur. Phys. J. C **67**, 425 (2010).
[39] E. Nakano, B. J. Schaefer, B. Stokic, B. Friman, and K. Redlich, Phys. Lett. B **682**, 401 (2010).
[40] J. Braun, H. -J. Pirner, K. Schwenzer, Phys. Rev. D **70**, 085016 (2004).
[41] M. Stephanov, Phys. Rev. D **73**, 094508 (2006).
[42] V. Skokov, K. Morita, and B. Friman, Phys. Rev. D **83**, 071502(R) (2011); B. Friman, Acta. Phys. Pol. B (Proc Suppl.) **5**, 707 (2012).
[43] P. Braun-Munzinger, K. Redlich, and J. Stachel, in *Quark-Gluon Plasma 3*, edited by R. C. Hwa and X. N. Wang (World Scientific, 2004).
[44] R. Hagedorn and K. Redlich, Z. Phys. C **27**, 541 (1985).
[45] A. Roberge and N. Weiss, Nucl. Phys. **B275**, 734 (1986).
[46] Y. Sakai, K. Kashiwa, H. Kouno, and M. Yahiro, Phys. Rev. D **77**, 051901(R) (2008).
[47] K. Morita, V. Skokov, B. Friman, and K. Redlich, Phys. Rev. D **84**, 076009 (2011).
[48] K. Morita, V. Skokov, B. Friman, and K. Redlich, Phys. Rev. D **84**, 074020 (2011).
[49] C. N. Yang and T. D. Lee, Phys. Rev. **87**, 404 (1952).
[50] M. Wagner, A. Walther, and B. -J. Schaefer, Comp. Phys. Comm. **181**, 756 (2010).
[51] K. Morita, talk given at The 4th Asian Triangle Heavy Ion Conference (ATHIC2012), November 16 2012, Pusan, Korea; K. Morita, B. Friman, K. Redlich, V. Skokov, in preparation.